

DR WP DE BEER

MBChB [UFS] M MED [SURG] CUM LAUDE

GENERAL SURGEON/ALGEMENE CHIRURG

SUITE 304
LINKSFIELD PARK CLINIC
24 12TH AVENUE
LINKSFIELD WEST
JOHANNESBURG

PHONE
011-4854741 [ROOMS]
011-2526322 [FAX]
Practice No: 0082295

Surgery for Pilonidal Sinus

This document will give you information about surgery for a pilonidal sinus. If you have any questions, you should ask your GP or other relevant health professional.

What is a pilonidal sinus?

A pilonidal sinus is a problem in the natal cleft (area just above the buttocks). Loose hairs fall off the neck or back and collect in the natal cleft, causing small holes to form in the skin or getting into existing holes. As the hairs carry bacteria, the holes can become infected. This causes an abscess to form or a discharge that is released through a tunnel (sinus) out onto the skin

What are the benefits of surgery?

Surgery is the most dependable way to remove the pilonidal sinus.

Are there any alternatives to surgery?

You can treat an occasional discharge with antibiotics.

What does the operation involve?

The operation is performed under a general anaesthetic. Your surgeon will remove the sinus and infected tissue.

What complications can happen?

1 General complications of any operation

- . Pain*
- . Bleeding*
- . Unsightly scarring*
- . Blood clots*

2 Specific complications of this operation

- . Partial breakdown of the wound*
- . Slow healing*
- . Numbness around the wound*

How soon will I recover?

*You should be able to go home the same day.
You should be able to return to work after two to three weeks.
Occasionally the pilonidal sinus comes back.*

Summary

Pilonidal sinus is a common problem in young people. It can cause an abscess or persistent discharge and is best treated by surgery.

NB!!! IF THE WARD DID NOT BOOK A FOLLOW-UP APPOINTMENT, PLEASE CALL THE ROOMS TO DO SO.